

## Solutions of APMO 2014

**Problem 1.** For a positive integer  $m$  denote by  $S(m)$  and  $P(m)$  the sum and product, respectively, of the digits of  $m$ . Show that for each positive integer  $n$ , there exist positive integers  $a_1, a_2, \dots, a_n$  satisfying the following conditions:

$$S(a_1) < S(a_2) < \dots < S(a_n) \text{ and } S(a_i) = P(a_{i+1}) \quad (i = 1, 2, \dots, n).$$

(We let  $a_{n+1} = a_1$ .) (*Problem Committee of the Japan Mathematical Olympiad Foundation*)

**Solution.** Let  $k$  be a sufficiently large positive integer. Choose for each  $i = 2, 3, \dots, n$ ,  $a_i$  to be a positive integer among whose digits the number 2 appears exactly  $k + i - 2$  times and the number 1 appears exactly  $2^{k+i-1} - 2(k + i - 2)$  times, and nothing else. Then, we have  $S(a_i) = 2^{k+i-1}$  and  $P(a_i) = 2^{k+i-2}$  for each  $i$ ,  $2 \leq i \leq n$ . Then, we let  $a_1$  be a positive integer among whose digits the number 2 appears exactly  $k + n - 1$  times and the number 1 appears exactly  $2^k - 2(k + n - 1)$  times, and nothing else. Then, we see that  $a_1$  satisfies  $S(a_1) = 2^k$  and  $P(a_1) = 2^{k+n-1}$ . Such a choice of  $a_1$  is possible if we take  $k$  to be large enough to satisfy  $2^k > 2(k + n - 1)$  and we see that the numbers  $a_1, \dots, a_n$  chosen this way satisfy the given requirements.

**Problem 2.** Let  $S = \{1, 2, \dots, 2014\}$ . For each non-empty subset  $T \subseteq S$ , one of its members is chosen as its *representative*. Find the number of ways to assign representatives to all non-empty subsets of  $S$  so that if a subset  $D \subseteq S$  is a disjoint union of non-empty subsets  $A, B, C \subseteq S$ , then the representative of  $D$  is also the representative of at least one of  $A, B, C$ . (*Warut Suksompong, Thailand*)

**Solution.** *Answer:*  $108 \cdot 2014!$ .

For any subset  $X$  let  $r(X)$  denotes the representative of  $X$ . Suppose that  $x_1 = r(S)$ . First, we prove the following fact:

$$\text{If } x_1 \in X \text{ and } X \subseteq S, \text{ then } x_1 = r(X).$$

If  $|X| \leq 2012$ , then we can write  $S$  as a disjoint union of  $X$  and two other subsets of  $S$ , which gives that  $x_1 = r(X)$ . If  $|X| = 2013$ , then let  $y \in X$  and  $y \neq x_1$ . We can write  $X$  as a disjoint union of  $\{x_1, y\}$  and two other subsets. We already proved that  $r(\{x_1, y\}) = x_1$  (since  $|\{x_1, y\}| = 2 < 2012$ ) and it follows that  $y \neq r(X)$  for every  $y \in X$  except  $x_1$ . We have proved the fact.

Note that this fact is true and can be proved similarly, if the ground set  $S$  would contain at least 5 elements.

There are 2014 ways to choose  $x_1 = r(S)$  and for  $x_1 \in X \subseteq S$  we have  $r(X) = x_1$ . Let  $S_1 = S \setminus \{x_1\}$ . Analogously, we can state that there are 2013 ways to choose  $x_2 = r(S_1)$  and for  $x_2 \in X \subseteq S_1$  we have  $r(X) = x_2$ . Proceeding similarly (or by induction), there are  $2014 \cdot 2013 \cdots 5$  ways to choose  $x_1, x_2, \dots, x_{2010} \in S$  so that for all  $i = 1, 2, \dots, 2010$ ,  $x_i = r(X)$  for each  $X \subseteq S \setminus \{x_1, \dots, x_{i-1}\}$  and  $x_i \in X$ .

We are now left with four elements  $Y = \{y_1, y_2, y_3, y_4\}$ . There are 4 ways to choose  $r(Y)$ . Suppose that  $y_1 = r(Y)$ . Then we clearly have  $y_1 = r(\{y_1, y_2\}) = r(\{y_1, y_3\}) = r(\{y_1, y_4\})$ . The only subsets whose representative has not been assigned yet are  $\{y_1, y_2, y_3\}$ ,  $\{y_1, y_2, y_4\}$ ,  $\{y_1, y_3, y_4\}$ ,  $\{y_2, y_3, y_4\}$ ,  $\{y_2, y_3\}$ ,  $\{y_2, y_4\}$ ,  $\{y_3, y_4\}$ . These subsets can be assigned in any way, hence giving  $3^4 \cdot 2^3$  more choices.

In conclusion, the total number of assignments is  $2014 \cdot 2013 \cdots 4 \cdot 3^4 \cdot 2^3 = 108 \cdot 2014!$ .

**Problem 3.** Find all positive integers  $n$  such that for any integer  $k$  there exists an integer  $a$  for which  $a^3 + a - k$  is divisible by  $n$ . (*Warut Suksompong, Thailand*)

**Solution.** *Answer:* All integers  $n = 3^b$ , where  $b$  is a nonnegative integer.

We are looking for integers  $n$  such that the set  $A = \{a^3 + a \mid a \in \mathbf{Z}\}$  is a complete residue system by modulo  $n$ . Let us call this property by (\*). It is not hard to see that  $n = 1$  satisfies (\*) and  $n = 2$  does not.

If  $a \equiv b \pmod{n}$ , then  $a^3 + a \equiv b^3 + b \pmod{n}$ . So  $n$  satisfies (\*) iff there are no  $a, b \in \{0, \dots, n-1\}$  with  $a \neq b$  and  $a^3 + a \equiv b^3 + b \pmod{n}$ .

First, let us prove that  $3^j$  satisfies (\*) for all  $j \geq 1$ . Suppose that  $a^3 + a \equiv b^3 + b \pmod{3^j}$  for  $a \neq b$ . Then  $(a-b)(a^2 + ab + b^2 + 1) \equiv 0 \pmod{3^j}$ . We can easily check mod 3 that  $a^2 + ab + b^2 + 1$  is not divisible by 3.

Next note that if  $A$  is not a complete residue system modulo integer  $r$ , then it is also not a complete residue system modulo any multiple of  $r$ . Hence it remains to prove that any prime  $p > 3$  does not satisfy (\*).

If  $p \equiv 1 \pmod{4}$ , there exists  $b$  such that  $b^2 \equiv -1 \pmod{p}$ . We then take  $a = 0$  to obtain the congruence  $a^3 + a \equiv b^3 + b \pmod{p}$ .

Suppose now that  $p \equiv 3 \pmod{4}$ . We will prove that there are integers  $a, b \not\equiv 0 \pmod{p}$  such that  $a^2 + ab + b^2 \equiv -1 \pmod{p}$ . Note that we may suppose that  $a \not\equiv b \pmod{p}$ , since otherwise if  $a \equiv b \pmod{p}$  satisfies  $a^2 + ab + b^2 + 1 \equiv 0 \pmod{p}$ , then  $(2a)^2 + (2a)(-a) + a^2 + 1 \equiv 0 \pmod{p}$  and  $2a \not\equiv -a \pmod{p}$ . Letting  $c$  be the inverse of  $b$  modulo  $p$  (i.e.  $bc \equiv 1 \pmod{p}$ ), the relation is equivalent to  $(ac)^2 + ac + 1 \equiv -c^2 \pmod{p}$ . Note that  $-c^2$  can take on the values of all non-quadratic residues modulo  $p$ . If we can find an integer  $x$  such that  $x^2 + x + 1$  is a non-quadratic residue modulo  $p$ , the values of  $a$  and  $c$  will follow immediately. Hence we focus on this latter task.

Note that if  $x, y \in \{0, \dots, p-1\} = B$ , then  $x^2 + x + 1 \equiv y^2 + y + 1 \pmod{p}$  iff  $p$  divides  $x + y + 1$ . We can deduce that  $x^2 + x + 1$  takes on  $(p+1)/2$  values as  $x$  varies in  $B$ . Since there are  $(p-1)/2$  non-quadratic residues modulo  $p$ , the  $(p+1)/2$  values that  $x^2 + x + 1$  take on must be 0 and all the quadratic residues.

Let  $C$  be the set of quadratic residues modulo  $p$  and 0, and let  $y \in C$ . Suppose that  $y \equiv z^2 \pmod{p}$  and let  $z \equiv 2w + 1 \pmod{p}$  (we can always choose such  $w$ ). Then  $y + 3 \equiv 4(w^2 + w + 1) \pmod{p}$ . From the previous paragraph, we know that  $4(w^2 + w + 1) \in C$ . This means that  $y \in C \implies y + 3 \in C$ . Unless  $p = 3$ , the relation implies that all elements of  $B$  are in  $C$ , a contradiction. This concludes the proof.

**Problem 4.** Let  $n$  and  $b$  be positive integers. We say  $n$  is  $b$ -discerning if there exists a set consisting of  $n$  different positive integers less than  $b$  that has no two different subsets  $U$  and  $V$  such that the sum of all elements in  $U$  equals the sum of all elements in  $V$ .

(a) Prove that 8 is a 100-discerning.

(b) Prove that 9 is not 100-discerning.

(*Senior Problems Committee of the Australian Mathematical Olympiad Committee*)

**Solution.**

(a) Take  $S = \{3, 6, 12, 24, 48, 95, 96, 97\}$ , i.e.

$$S = \{3 \cdot 2^k : 0 \leq k \leq 5\} \cup \{3 \cdot 2^5 - 1, 3 \cdot 2^5 + 1\}.$$

As  $k$  ranges between 0 to 5, the sums obtained from the numbers  $3 \cdot 2^k$  are  $3t$ , where  $1 \leq t \leq 63$ . These are 63 numbers that are divisible by 3 and are at most  $3 \cdot 63 = 189$ .

Sums of elements of  $S$  are also the numbers  $95 + 97 = 192$  and all the numbers that are sums of 192 and sums obtained from the numbers  $3 \cdot 2^k$  with  $0 \leq k \leq 5$ . These are 64 numbers that are all divisible by 3 and at least equal to 192. In addition, sums of elements of  $S$  are the numbers 95 and all the numbers that are sums of 95 and sums obtained from the numbers  $3 \cdot 2^k$  with  $0 \leq k \leq 5$ . These are 64 numbers that are all congruent to  $-1 \pmod{3}$ .

Finally, sums of elements of  $S$  are the numbers 97 and all the numbers that are sums of 97 and sums obtained from the numbers  $3 \cdot 2^k$  with  $0 \leq k \leq 5$ . These are 64 numbers that are all congruent to  $1 \pmod{3}$ .

Hence there are at least  $63 + 64 + 64 + 64 = 255$  different sums from elements of  $S$ . On the other hand,  $S$  has  $2^8 - 1 = 255$  non-empty subsets. Therefore  $S$  has no two different subsets with equal sums of elements. Therefore, 8 is 100-discerning.

(b) Suppose that 9 is 100-discerning. Then there is a set  $S = \{s_1, \dots, s_9\}$ ,  $s_i < 100$  that has no two different subsets with equal sums of elements. Assume that  $0 < s_1 < \dots < s_9 < 100$ .

Let  $X$  be the set of all subsets of  $S$  having at least 3 and at most 6 elements and let  $Y$  be the set of all subsets of  $S$  having exactly 2 or 3 or 4 elements greater than  $s_3$ .

The set  $X$  consists of

$$\binom{9}{3} + \binom{9}{4} + \binom{9}{5} + \binom{9}{6} = 84 + 126 + 126 + 84 = 420$$

subsets of  $S$ . The set in  $X$  with the largest sums of elements is  $\{s_4, \dots, s_9\}$  and the smallest sums is in  $\{s_1, s_2, s_3\}$ . Thus the sum of the elements of each of the 420 sets in  $X$  is at least  $s_1 + s_2 + s_3$  and at most  $s_4 + \dots + s_9$ , which is one of  $(s_4 + \dots + s_9) - (s_1 + s_2 + s_3) + 1$  integers. From the pigeonhole principle it follows that  $(s_4 + \dots + s_9) - (s_1 + s_2 + s_3) + 1 \geq 420$ , i.e.,

$$(s_4 + \dots + s_9) - (s_1 + s_2 + s_3) \geq 419. \quad (1)$$

Now let us calculate the number of subsets in  $Y$ . Observe that  $\{s_4, \dots, s_9\}$  has  $\binom{6}{2}$  2-element subsets,  $\binom{6}{3}$  3-element subsets and  $\binom{6}{4}$  4-element subsets, while  $\{s_1, s_2, s_3\}$  has exactly 8 subsets. Hence the number of subsets of  $S$  in  $Y$  equals

$$8 \left( \binom{6}{2} + \binom{6}{3} + \binom{6}{4} \right) = 8(15 + 20 + 15) = 400.$$


The set in  $Y$  with the largest sum of elements is  $\{s_1, s_2, s_3, s_6, s_7, s_8, s_9\}$  and the smallest sum is in  $\{s_4, s_5\}$ . Again, by the pigeonhole principle it follows that  $(s_1 + s_2 + s_3 + s_6 + s_7 + s_8 + s_9) - (s_4 + s_5) + 1 \geq 400$ , i.e.,

$$(s_1 + s_2 + s_3 + s_6 + s_7 + s_8 + s_9) - (s_4 + s_5) \geq 399. \quad (2)$$

Adding (1) and (2) yields  $2(s_6 + s_7 + s_8 + s_9) \geq 818$ , so that  $s_9 + 98 + 97 + 96 \geq s_9 + s_8 + s_7 + s_6 \geq 409$ , i.e.  $s_9 \geq 118$ , a contradiction with  $s_9 < 100$ . Therefore, 9 is not 100-discerning.

**Problem 5.** Circles  $\omega$  and  $\Omega$  meet at points  $A$  and  $B$ . Let  $M$  be the midpoint of the arc  $AB$  of circle  $\omega$  ( $M$  lies inside  $\Omega$ ). A chord  $MP$  of circle  $\omega$  intersects  $\Omega$  at  $Q$  ( $Q$  lies inside  $\omega$ ). Let  $\ell_P$  be the tangent line to  $\omega$  at  $P$ , and let  $\ell_Q$  be the tangent line to  $\Omega$  at  $Q$ . Prove that the circumcircle of the triangle formed by the lines  $\ell_P$ ,  $\ell_Q$ , and  $AB$  is tangent to  $\Omega$ . (*Ilya Bogdanov, Russia and Medeubek Kungozhin, Kazakhstan*)

**Solution.** Denote  $X = AB \cap \ell_P$ ,  $Y = AB \cap \ell_Q$ , and  $Z = \ell_P \cap \ell_Q$ . Without loss of generality we have  $AX < BX$ . Let  $F = MP \cap AB$ .



Denote by  $R$  the second point of intersection of  $PQ$  and  $\Omega$ ; by  $S$  the point of  $\Omega$  such that  $SR \parallel AB$ ; and by  $T$  the point of  $\Omega$  such that  $RT \parallel \ell_P$ . Since  $M$  is the midpoint of arc  $AB$ , the tangent  $\ell_M$  at  $M$  to  $\omega$  is parallel to  $AB$ , so  $\angle(AB, PM) = \angle(PM, \ell_P)$ . Therefore we have  $\angle PRT = \angle MPX = \angle PFX = \angle PRS$ . Thus the point  $Q$  is the midpoint of the arc  $TQS$  of  $\Omega$ , hence  $ST \parallel \ell_Q$ . So the corresponding sides of the triangles  $RST$  and  $XYZ$  are parallel, and there exist a homothety  $h$  mapping  $RST$  to  $XYZ$ .

Let  $D$  be the second point of intersection of  $XR$  and  $\Omega$ . We claim that  $D$  is the center of the homothety  $h$ ; since  $D \in \Omega$ , this implies that the circumcircles of triangles  $RST$  and  $XYZ$  are tangent, as required. So, it remains to prove this claim. In order to do this, it suffices to show that  $D \in SY$ .

By  $\angle PFX = \angle XPF$  we have  $XF^2 = XP^2 = XA \cdot XB = XD \cdot XR$ . Therefore,  $\frac{XF}{XD} = \frac{XR}{XF}$ , so the triangles  $XDF$  and  $XFR$  are similar, hence  $\angle DFX = \angle XRF = \angle DRQ = \angle DQY$ ; thus the points  $D, Y, Q,$  and  $F$  are concyclic. It follows that  $\angle YDQ = \angle YFQ = \angle SRQ = 180^\circ - \angle SDQ$  which means exactly that the points  $Y, D,$  and  $S$  are collinear, with  $D$  between  $S$  and  $Y$ .